



Fate, Behavior & Modeling of Spilled Oil Sands Products (Freshwater & Marine Environments)

Bruce Hollebone ESTS, Environment Canada

Oil Sands Products Training Seattle, WA 16 April 2013



Oil Fate, Behaviour and Models

Model:

Quantitative or semi-quantitative prediction of oil movement in environment (trajectory), and transformation by environmental factors (weathering)

Oil Behaviour:

A physical/chemical transformation of the oil, a "weathering" process

Oil Fate:

The eventual end-point state an oil may reach






Weathering

- In the environment crude oil and refined products are modified by natural physical, chemical, and biological processes that change their composition and fate.
- These processes are collectively referred to as weathering
- Primary processes include: evaporation, formation of emulsions, dissolution, photochemical and biological oxidation.
- The oil properties and how they change in the environment are one of the major drivers of oil behaviour (what it does), fate (where it goes), persistance (how long it lasts) and effects (who gets hurt).





Behaviour and Fate



Selected Physical Properties


Commercial Type		Heavy Sour Synbit	Sweet Synthetic Crude	Condensate	Heavy Sour Dilsynbit	Heavy Sour Dilbit	Light Sweet Crude
Oil		MacKay River Bitumen Diluted with Light Synthetic Crude Oil	Syncrude Synthetic Light Crude Oil	CRW Condensate	Albian Heavy Synthetic	Wabasca Heavy	West Texas Intermediate
Density (g/mL) 0 ºC					0.9463	1.0109	0.8594
	15 ºC	0.943	0.873	0.734	0.9372	0.9572	0.8474
Viscosity (mPa∙s)	0 ºC				465	1007933	19.2
	15 ºC	241.9	6	0.6	156	128100	8.6
Flash Point	(ºC)	10	<-5	<-5	-23	151	-10
Pour Point	(ºC)	<-24	<-24	<-25	<-30	-6	-22
Sulfur	(%)				2.3	4.7	0.90%







Chemistries


Saturate n-Alkanes









Polycyclic Aromatic Hydrocarbons (PAH)









ent Environnement Canada



Effects of Weathering on Models

- Weathering affects oil characteristics, density, viscosity, surface/interfacial tensions
- Physical and chemical changes cause differences in trajectory behaviours and oil fate

Oil Type	Wind Drift or Leeway (%)		
Gasoline	34		
Diesel	34		
Fresh IFO	3		
Fresh Crude	34		
Weathered IFO	23		
Emulsified Oil	12		
Scattered Tarballs	0.52		






Factors: Evaporation

- Best understood weathering process.
- Evaporation is influential behaviour component
- A light crude can lose as much as 75% in two days and a medium crude as much as 40%
- A physical/thermodynamic process, molecules are removed from the oil, but not changed chemically.
- One of the best studied types of weathering, many major types of models for oil -> air evaporation.






Evaporation of Oil Sands Products



Dissolution and Solubility

- Not a major effect on oil behaviour, but can have large effects on organisms and habitats
- Affected by oil concentration, mixing energy, temperature and time
- Can be limited by diffusion rates through Oil
- Physical models well developed





Factors: Photo-oxidation

•Observed to cause increase in density in real-world spills

- •"Skin" formation
- •Appears to increase oxidized components of oil


•Enhances water uptake and emulsion formation

•No models for behaviour









Environ Canada

anment

Environnement Canada




Factors: Water Uptake

- Formation of finely dispersed water droplets in an oil phase
- Emulsion densities increase, viscosities greatly increase
- Changes state of oil, but not chemistry.
- Changes transport: oil is lower in water, has higher stickiness and much higher viscosity
- "Entrained water"— simple viscosity-bound water in oil may persist for significant amounts of time
- Models for water uptake and retention are not well developed --- all data is empirical/experimental










Environnement Ceneda



Evaporation and Photo-oxidation Effects on Water Content of Oil








Factors: Sediment Interaction

- Observed sedimentation of sunken oil at several spills
 - Quantitative data from spills of opportunity is sparse.
- Quantitative models just now starting to be developed
- Several modes of uptake
 - Suspended particulates (OMA/OFI/OSA/SPM, etc...)
 - Surf-zone interactions












ent Environneme Canada








Dispersion and Sediment Interaction

- Dispersion of oil into small droplets occurs in highenergy environments: breaking waves, storms, floods, surf zones.
- Dispersion can promote rapid weathering because of proportionately large surface areas to volume of oil
 - Dissolution
 - Emulsification
 - Sediment uptake
 - Water uptake
- Sinking and "tarball formation"
- Models are under active development



Khelifa et al, 2008 **Canadă**



Water Surface





Envire

t Environnement Canada



Immediately After Agitation





t Environnement Canada



A Few Minutes Later





t Environnement Canada




Factors: Temperature


- Oil density increases with decreasing temperature
- Cyclic daily and seasonal variations, with additional weather factors
- Temperature affects many oil properties (viscosity, interfacial tensions)
 - Potential for non-linear interactions
- Empirically well understood, practical models exist







Sediment Uptake vs Temperature







Chemical Weathering

• Chemical distribution of oil changes in the environment.

Removal:

- Evaporation
- Dissolution and dispersal

Transformational:


- Chemical degradation/oxidation (eg photo-enhanced)
- Biological degradation





Weathering changes to oil

- Largest changes occur rapidly after spill, with slow degradation later
- Rapid removal of light ends
 <C25 (F1 + F2)
- The fate of the oil strongly affects degredation
- Densities uniformly increased with exposure
- Oil does not weather evenly; highly degraded skins can contain less weathered oil.
- Models are mostly empirical







Effects of Evaporation

Polycyclic Aromatic Hydrocarbons (PAH) Saturate n-Alkanes Albian Heavy Synthetic, 22.6% weathered Albian Heavy Synthetic 22.6% weathered 260 200 160 100 60 2.5 1000 2.0 800 Conc. (mg/g) Conc. (µg/g oil) 1.5 600 5 2 5 5 E C C 1.0 400 200 0.5 0.0 C2-N C3-N 04-N S0-N CI-N G-D-D 5 2-1-2 с В 4 0-0-0 Wabisca Heavy, 10.7% weathered 15 Wabisca Heavy, 10.7% weathered 15.0 10 500 12.0 400 Conc. (µg/g oil) Conc. (mg/g) 9.0 300 6.0 200 3.0 100 0.0 Π C2-N 3-N 04-N G-P <u>д</u>-C2-P C3-P С4-Р 8 2-00 01-N West Texas Intermediate 150 West Texas Intermediate 31.7% w 31.7% w 75 -0 -율 10 2000 8 1500 6 4 1000 2 500 0 n-C11 n-C14 Phytane n-C24 n-C30 n-C33 n-C36 n-C39 n-C8 n-C17 n-C21 n-C27 0

C1-D C2-D 3 5 5 C2-F 5 0-00 5 C2-C Other EPA Priority PA-is





Environnement Canada



Other EPA Priority PAHs

3 3


Biodegredation

- Long-term action of microbes on oil
- Almost reverse of photo-oxidation: smaller molecules affected first, straight-chain alkanes preferred, followed by unalkylated aromatics.
- Can last for months to years.
- Aerobic can be much faster than anaerobic.
- Availability of fertilizers and electron acceptors






Biodegradation of Heavy Oil



Bio-oxidation in natural samples



Open Questions

- Changes to dilbit chemistry by evaporation of diluent components (rate of change)
- Dispersion of oil in water (droplet size, rise time, coalescence)
- Resuspension and remobilization potential
- Dissolution in water & bioavailability
- Toxicity, chemical and physical
- Persistence
- Interactions of factors to affect behaviours





Questions? Comments?





